

CUSHING'S DISEASE HAS  
HELD YOU DOWN LONG ENOUGH

## START A NEW CHAPTER WITH ISTURISA



Learn about how you may benefit from treatment

Model shown is not an actual patient.

### Indications and Usage

ISTURISA® (osilodrostat) is used to treat adults with Cushing's disease who cannot have pituitary surgery, or who have had pituitary surgery, but the surgery did not cure their Cushing's disease.

### Important Safety Information

**Hypocortisolism:** Treatment with ISTURISA may cause symptoms associated with low levels of cortisol in your blood (hypocortisolism). Tell your healthcare provider right away if you experience more than one of the following symptoms, as these may be symptoms of very low cortisol level, known as adrenal insufficiency: nausea, vomiting, tiredness (fatigue), low blood pressure, stomach (abdominal) pain, loss of appetite, dizziness. If you get symptoms of hypocortisolism while taking ISTURISA, your healthcare provider may change your dose or ask you to stop taking it.

**Please see Important Safety Information throughout and accompanying Full Prescribing Information.**



# ISTURISA HELPS BRING CORTISOL LEVELS BACK TO NORMAL<sup>1</sup>

ISTURISA (is tur ee' sah) is a prescription medicine used to treat adult patients with Cushing's disease who can't have pituitary surgery, or who have had pituitary surgery but were not cured of their Cushing's disease.<sup>1</sup>

**ISTURISA can help you reach and maintain normal cortisol levels.<sup>1</sup>**



Model shown is not an actual patient.

- The symptoms of Cushing's disease are caused by too much cortisol. The way to relieve them is to bring cortisol levels back to normal<sup>2</sup>
- ISTURISA works to reduce the amount of cortisol made in your adrenal glands<sup>1</sup>
- With ISTURISA, less cortisol is made, so less cortisol is released into your bloodstream

## **Important Safety Information**

**Heart Problem or Heart Rhythm Problem:** ISTURISA may cause an irregular heartbeat which could be a sign of a heart problem called QT prolongation. Call your healthcare provider right away if you have irregular heartbeats.

**Please see Important Safety Information throughout and the accompanying Full Prescribing Information**



## START A NEW CHAPTER

We'll be  
here to support  
you every step  
of the way

### Isturisa (osilodrostat) offers:

- Proven results in patients with Cushing's disease<sup>1</sup>
- Dosing that can be individualized to your specific needs<sup>1</sup>
- Tablets that can be taken with or without food<sup>1</sup>
- Dedicated nurse, pharmacist, and insurance specialist support through the R.A.R.E. program

### Personalized treatment for you

ISTURISA comes in 3 different tablet strengths: 1 mg, 5 mg and 10 mg.<sup>1</sup> You may take a combination of different tablets based on how you respond.<sup>1</sup> This allows your doctor to determine the precise dose of ISTURISA that works best for you.

In the clinical study for ISTURISA, the most common side effects were<sup>1</sup>:

- Very low cortisol levels (hypocortisolism), which includes flu-like symptoms
- Tiredness (fatigue)
- Nausea
- Headache
- Swelling of the legs, ankles, or other signs of fluid retention (edema)



### Important Safety Information

**Increase in Other Adrenal Hormone Levels:** Your other adrenal hormones may increase when you take ISTURISA. Your healthcare provider may monitor you for the symptoms associated with these hormonal changes while you are taking ISTURISA such as low potassium (hypokalemia), high blood pressure (hypertension), swelling (edema) in the legs, ankles, or other signs of fluid retention, excessive facial or body hair growth (hirsutism), acne (in women). Call your healthcare provider if you have any of these side effects.

# CORTISOL AND CUSHING'S DISEASE

Life with Cushing's disease isn't easy. For many, it has been a long road. Cushing's disease is caused by a non-cancerous pituitary tumor that signals your adrenal glands to constantly produce cortisol. Your body needs cortisol for many processes, including your ability to maintain normal blood pressure and heart function, process carbohydrates, fats, and proteins – even your ability to fight infection.<sup>3</sup>

**Too much cortisol can lead to obesity, diabetes, high cholesterol levels, high blood pressure, muscle and bone weakness, excess hair growth, increased bruising, depression, anxiety, and frequent infections.<sup>3</sup>**



**In a clinical study, Isturisa® (osilodrostat) was proven to lower cortisol levels<sup>1</sup>**

COMPLETE RESPONSE

3x

3x as many patients had a complete response\* to ISTURISA compared to placebo

2/3 OF PATIENTS



2/3 of patients on ISTURISA had normal cortisol levels after 48 weeks

\*A complete response was defined as patients who had normal cortisol levels after week 34, and who neither discontinued taking ISTURISA or increased their dose after week 26.

## Important Safety Information

**Most common side effects include** very low cortisol levels (adrenal insufficiency), tiredness (fatigue), nausea, headache, swelling of the legs, ankles or other signs of fluid retention (edema). These are not all of the possible side effects of ISTURISA.

**To report SUSPECTED SIDE EFFECTS, contact Recordati Rare Diseases Inc. at 1-888-575-8344, or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

# R.A.R.E.<sup>TM</sup>

Recordati Access, Resources, and Engagement



**INSURANCE SPECIALISTS**  
to help get you the best  
coverage for your medications



**PHARMACISTS**  
who are available to you  
24/7, every day of the year



**A NURSE AMBASSADOR**  
to answer your questions and  
support you during treatment

**R.A.R.E. is a patient support program designed to make it easier for you to access and stay on your treatment**

Once you're prescribed Isturisa<sup>®</sup> (osilodrostat) you are automatically enrolled in the R.A.R.E. program.

Our knowledgeable and reliable professionals have been fully trained about ISTURISA and are dedicated to supporting your treatment goals.

You'll have a dedicated nurse and 24/7 pharmacist support, financial assistance,\* education, and more.

\*with the cost of ISTURISA

MEDICATION  
FOR NO MORE THAN  
**\$20**  
PER MONTH<sup>†</sup>

**R.A.R.E. can help eligible commercially insured patients get their medication for no more than \$20 per month.**

For patients who are not eligible for copay support or who need additional financial assistance, R.A.R.E. may be able to connect you with alternative forms of financial support.<sup>†</sup>

<sup>†</sup>Restrictions, limitations, and/or eligibility requirements apply

**Please see Important Safety Information throughout and the accompanying Full Prescribing Information.**

CUSHING'S DISEASE HAS HELD YOU DOWN LONG ENOUGH

# IT'S TIME TO START A NEW CHAPTER WITH ISTURISA

Learn more at [www.isturisa.com](http://www.isturisa.com)

## ISTURISA<sup>®</sup> (osilodrostat) Important Safety Information

### Indications and Usage

ISTURISA (osilodrostat) is used to treat adults with Cushing's disease who cannot have pituitary surgery, or who have had pituitary surgery, but the surgery did not cure their Cushing's disease.

### Important Safety Information:

- **Hypocortisolism:** Treatment with ISTURISA may cause symptoms associated with low levels of cortisol in your blood (hypocortisolism). Tell your healthcare provider right away if you experience more than one of the following symptoms, as these may be symptoms of very low cortisol level, known as adrenal insufficiency: nausea, vomiting, tiredness (fatigue), low blood pressure, stomach (abdominal) pain, loss of appetite, dizziness. If you get symptoms of hypocortisolism while taking ISTURISA, your healthcare provider may change your dose or ask you to stop taking it.
- **Heart Problem or Heart Rhythm Problem:** ISTURISA may cause an irregular heartbeat which could be a sign of a heart problem called QT prolongation. Call your healthcare provider right away if you have irregular heartbeats.
- **Increase in Other Adrenal Hormone Levels:** Your other adrenal hormones may increase when you take ISTURISA. Your healthcare provider may monitor you for the symptoms associated with these hormonal changes while you are taking ISTURISA such as low potassium (hypokalemia), high blood pressure (hypertension), swelling (edema) in the legs, ankles, or other signs of fluid retention, excessive facial or body hair growth (hirsutism), acne (in women). Call your healthcare provider if you have any of these side effects.
- **Most common side effects include** very low cortisol levels (adrenal insufficiency), tiredness (fatigue), nausea, headache, swelling of the legs, ankles or other signs of fluid retention (edema). These are not all of the possible side effects of ISTURISA.

**To report SUSPECTED SIDE EFFECTS, contact Recordati Rare Diseases Inc. at 1-888-575-8344, or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

**Before taking ISTURISA, tell your doctor about all of your medical conditions, including if you:**

- have or had heart problems, such as an irregular heartbeat, including a condition called prolonged QT syndrome (QT internal prolongation). Your healthcare provider will check the electrical signal of your heart (called an electrocardiogram) before you start taking ISTURISA, 1 week after starting ISTURISA, and as needed after that.
- have a history of low levels of potassium or magnesium in your blood.
- have liver problems.
- are breastfeeding or plan to breastfeed. It is not known if ISTURISA passes into your breast milk. **You should not** breastfeed if you take ISTURISA and for 1 week after stopping treatment.

**Tell your healthcare provider about all the medicines you take, including** prescription medicines, over-the-counter medicines, vitamins, and herbal supplements. Especially tell your healthcare provider if you take medicines used to treat certain heart problems. Ask your healthcare provider if you are not sure whether your medicine is used to treat heart problems.